

A Necessary and Sufficient Condition for Invertibility of Adapted Perturbations of Identity on Wiener Space*

A. S. Üstünel

Abstract: Let (W, H, μ) be the classical Wiener space, assume that $U = I_W + u$ is an adapted perturbation of identity satisfying the Girsanov identity. Then, U is invertible if and only if the kinetic energy of u is equal to the relative entropy of the measure induced with the action of U on the Wiener measure μ , in other words U is invertible if and only if

$$\frac{1}{2} \int_W |u|_H^2 d\mu = \int_W \frac{dU\mu}{d\mu} \log \frac{dU\mu}{d\mu} d\mu.$$

Une condition nécessaire et suffisante pour l'inversibilité de perturbations d'identité adaptées sur l'espace de Wiener

Resumé: Soit (W, H, μ) l'espace de Wiener classique, et soit $U = I_W + u$ une perturbation d'identité adaptée satisfaisant à l'identité de Girsanov. Alors U est inversible si et seulement si l'énergie cinétique de u est égale à l'entropie relative de la mesure induite par l'action de U sur la mesure de Wiener μ ; en d'autres termes U est inversible si et seulement si

$$\frac{1}{2} \int_W |u|_H^2 d\mu = \int_W \frac{dU\mu}{d\mu} \log \frac{dU\mu}{d\mu} d\mu.$$

*To appear in Comptes Rendus Mathématiques, Vol. 346

1 Version française abrégée

Soit (W, H, μ) l'espace de Wiener classique: $W = C([0, 1], \mathbb{R}^d)$, $d \geq 1$ μ est la mesure gaussienne standard et H est l'espace de Cameron-Martin dont le produit scalaire et la norme sont notés respectivement par $(h, k)_H = \int_0^1 \dot{h}_s \cdot \dot{k}_s ds$ et par $|\cdot|_H$. Nous noterons par $(\mathcal{F}_t, t \in [0, 1])$ la filtration canonique du mouvement brownien canonique, éventuellement complétée. Soit maintenant $u \in L_a^0(\mu, H)$, où cette dernière représente les classes d'équivalences de variables aléatoires à valeurs dans H telles que leurs dérivées temporelles sont adaptées à la filtration brownienne. On notera par $\rho(u)$ l'exponentielle de Girsanov définie par

$$\rho(u) = \exp \left(\int_0^1 \dot{u}_s \cdot dW_s - \frac{1}{2} \int_0^1 |\dot{u}_s|^2 ds \right)$$

Nous avons le résultat suivant:

Theorem 1 *Supposons que $E[\rho(-u)] = 1$ et notons par U l'application $I_W + u$. Alors les propriétés suivantes sont équivalentes:*

1. *L'application U est p.s. inversible, son inverse peut s'écrire comme $V = I_W + v$ avec $v \in L_a^0(\mu, H)$,*
2. *L'équation différentielle stochastique suivante*

$$\begin{aligned} dV_t &= -\dot{u}_t \circ V dt + dW_t \\ V_0 &= 0 \end{aligned}$$

possède une solution (forte) unique,

3. *On a l'identité*

$$\frac{1}{2} \int_W |u|_H^2 d\mu = \int_W \frac{dU\mu}{d\mu} \log \frac{dU\mu}{d\mu} d\mu.$$

L'équivalence entre (1) et (2) a déjà démontrée dans [9] et le reste de cet article sera consacré à la preuve de l'équivalence entre (1) et (3).

2 Main results

Let (W, H, μ) the classical Wiener space on $\mathbb{R}^d$, we begin with the following proposition whose proof follows from the Girsanov theorem

Proposition 1 *Assume that $L = \rho(-v)$, where $v \in L_a^0(\mu, H)$, i.e., v is adapted and $\int_0^1 |\dot{v}_s|^2 ds < \infty$ a.s. Then there exists $U = I_W + u$, with $u : W \rightarrow H$ adapted such that $U\mu = L\mu$ and $E[\rho(-u)] = 1$ if and only if the following condition is satisfied:*

$$1 = L_t \circ U E[\rho(-u^t)|\mathcal{U}_t] \quad (2.1)$$

$$= L_t \circ U E[\rho(-u)|\mathcal{U}_t] \quad (2.2)$$

almost surely for any $t \in [0, 1]$, where u^t is defined as $u^t(\tau) = \int_0^{t \wedge \tau} \dot{u}_s ds$ and $\mathcal{U}_t$ is the sigma algebra generated by $(w(\tau) + u(\tau), \tau \leq t)$.

Let us calculate $E[\rho(-u^t)|\mathcal{U}_t] = E[\rho(-u)|\mathcal{U}_t]$ in terms of the innovation process associated to U . Recall that the term innovation, which originates from the filtering theory is defined as (cf.[3] and [7])

$$Z_t = U_t - \int_0^t E[\dot{u}_s|\mathcal{U}_s] ds$$

and it is a μ -Brownian motion with respect to the filtration $(\mathcal{U}_t, t \in [0, 1])$. A similar proof as the one in [3] shows that any martingale with respect to the filtration of U can be represented as a stochastic integral with respect to Z . Hence, by the positivity assumption, $E[\rho(-u)|\mathcal{U}_t]$ can be written as an exponential martingale

$$E[\rho(-u)|\mathcal{U}_t] = \exp \left(- \int_0^t (\dot{\xi}_s, dZ_s) - \frac{1}{2} \int_0^t |\dot{\xi}_s|^2 ds \right).$$

A double utilization of the Girsanov theorem gives the following explicit result:

Proposition 2 *We have*

$$E[\rho(-\delta u)|\mathcal{U}] = \exp \left(- \int_0^1 (E[\dot{u}_s|\mathcal{U}_s], dZ_s) - \frac{1}{2} \int_0^1 |E[\dot{u}_s|\mathcal{U}_s]|^2 ds \right), \quad (2.3)$$

and

$$E[\rho(-\delta u)|\mathcal{U}_t] = \exp \left(- \int_0^t (E[\dot{u}_s|\mathcal{U}_s], dZ_s) - \frac{1}{2} \int_0^t |E[\dot{u}_s|\mathcal{U}_s]|^2 ds \right), \quad (2.4)$$

almost surely.

Combining Propositions 1 and 2, we obtain

Theorem 2 *A necessary and sufficient condition for the relation (2.1) is that*

$$E[\dot{u}_t|\mathcal{U}_t] = -\dot{v}_t \circ U$$

dt × dμ-almost surely.

Now we state and prove the main theorem (cf. also [2] for related problems):

Theorem 3 *Assume that $u \in L^2(\mu, H) \cap L_a^0(\mu, H)$ with $E[\rho(-u)] = 1$. Define L as*

$$L = \frac{dU\mu}{d\mu} = \rho(-v)$$

where $v \in L_a^0(\mu, H)$ is given by the Itô representation theorem. The map $U = I_W + u$ is then almost surely invertible with its inverse $V = I_W + v$ if and only if

$$2E[L \log L] = E[|u|_H^2].$$

In other words, U is invertible if and only if

$$H(U\mu|\mu) = \frac{1}{2} \|u\|_{L^2(\mu, H)}^2,$$

where $H(U\mu|\mu)$ denotes the entropy of $U\mu$ with respect to μ .

Proof: Since U represents $Ld\mu$, we have, from Theorem 2, $E[\dot{u}_s|\mathcal{U}_s] + \dot{v}_s \circ U = 0$ $ds \times d\mu$ -almost surely. Hence, from the Jensen inequality $E[|v \circ U|_H^2] \leq E[|u|_H^2]$. Moreover the Girsanov theorem gives

$$2E[L \log L] = E[|v|_H^2 L] = E[|v \circ U|_H^2] = E\left[\int_0^1 |E[\dot{u}_s|\mathcal{U}_s]|^2 ds\right].$$

Hence the hypothesis implies that

$$E[|u|_H^2] = E\left[\int_0^1 |E[\dot{u}_s|\mathcal{U}_s]|^2 ds\right].$$

From which we deduce that $\dot{u}_s = E[\dot{u}_s|\mathcal{U}_s]$ $ds \times d\mu$ -almost surely. Finally we get $\dot{u}_s + \dot{v}_s \circ U = 0$ $ds \times d\mu$, which is a necessary and sufficient condition for the invertibility of U (cf.[7, 8, 9]) . The necessity is obvious.

□

Corollary 1 *With the notations of theorem, U is not invertible if and only if we have*

$$\frac{1}{2} E[|u|_H^2] > H(U\mu|\mu).$$

Remark: This result gives an enlightenment about the celebrated counter example of Tsirelson, cf. [4].

Acknowledgement: Some parts of this work has been done while the author was visiting the Departement of Mathematics of Bilkent University, Ankara, Turkey.

References

- [1] D. Allinger and S.K. Mitter: “ New results on the innovations problem for nonlinear filtering”. Stochastics **4**, no. 4, 339–348, 1980.
- [2] D. Feyel, A.S. Üstünel and M. Zakai: “Realization of Positive Random Variables via Absolutely Continuous Transformations of Measure on Wiener Space”. Probability Surveys, Vol. 3, (electronic) p.170-205, 2006.
- [3] M. Fujisaki, G. Kallianpur and H. Kunita: “Stochastic differential equations for the non linear filtering problem”. Osaka J. Math., **9**, p. 19-40, 1972.

- [4] N. Ikeda and S. Watanabe: *Stochastic Differential Equations and Diffusion Processes*. North Holland, Amsterdam (Kodansha Ltd., Tokyo), 1981.
- [5] A. S. Üstünel: *Introduction to Analysis on Wiener Space*. Lecture Notes in Math. Vol. 1610. Springer, 1995.
- [6] A. S. Üstünel: *Analysis on Wiener Space and Applications*. Electronic text at the site <http://www.finance-research.net/>.
- [7] A. S. Üstünel and M. Zakai: *Transformation of Measure on Wiener Space*. Springer Verlag, 1999.
- [8] A. S. Üstünel and M. Zakai: “The invertibility of adapted perturbations of identity on the Wiener space”. C. R. Acad. Sci. Paris, Série I, **342**, p. 689-692, 2006.
- [9] A. S. Üstünel and M. Zakai: “Sufficient conditions for the invertibility of adapted perturbations of identity on the Wiener space”. Probab. Theory Relat. Fields, **139**, p. 207-234, 2007.

A. S. Üstünel, Telecom-Paristech, Dépt. Infres,
 46, rue Barrault, 75013, Paris, France
 ustunel@enst.fr